

Interventional Radiology: Diagnostic Angiogram



Name: _____ Date: _____ Time: _____ Buck Pavilion 5



Please arrange a ride home from the procedure



If you are currently taking Coumadin ask your doctor for instructions to stop.

5 days before	3 days before	2 days before	Procedure Day
<p data-bbox="283 552 625 673">  Begin arranging a ride home from your procedure. </p> <p data-bbox="409 803 546 933">  </p> <p data-bbox="283 950 646 1242"> If you use Coumadin please call your physician or your anticoagulation clinic nurse for instructions to stop your Coumadin. If you have not received these instructions, please call the Radiology Clinic RN at 206 341-0152. </p>	<p data-bbox="808 560 945 690">  </p> <p data-bbox="693 706 1018 885"> If you are Allergic to IV CONTRAST Call the Radiology Clinic RN for instructions at (206) 341-0152. </p> <p data-bbox="808 901 945 1031">  </p> <p data-bbox="724 1015 976 1079"> If you use Coumadin STOP TODAY </p>	<p data-bbox="1207 544 1344 673">  </p> <p data-bbox="1081 673 1470 950"> You will receive a call today from the Radiology Clinic RN who will go over your instructions. If you have questions please ask at that time. If you live outside of Seattle please call and leave a number you can be reached at. </p> <p data-bbox="1134 1047 1333 1079"> NIGHT BEFORE </p> <p data-bbox="1081 1104 1459 1380">  If you are allergic to Contrast you should now have a Prescription for Prednisone and Benadryl. Take one Prednisone tablet at 4PM and 10 PM the day before your procedure. </p>	<p data-bbox="1501 544 1606 665">  </p> <p data-bbox="1617 568 1837 673"> Take your morning medications with sips of water. </p> <p data-bbox="1491 706 1606 820">  </p> <p data-bbox="1617 706 1879 885"> No solid food 6 hours before your procedure. You may drink clear liquids up until 2 hours before your procedure. </p> <p data-bbox="1491 901 1606 1006">  </p> <p data-bbox="1617 917 1848 1079"> If you are a diabetic only take half of your insulin dose and if you take diabetic pills hold them that morning. </p> <p data-bbox="1501 1112 1606 1209">  </p> <p data-bbox="1617 1112 1816 1209"> please allow 4-6 hours for your recovery. </p> <p data-bbox="1491 1258 1606 1364">  </p> <p data-bbox="1617 1250 1869 1404"> If you are allergic to Contrast take your remaining pills one hour before your procedure. </p>
